

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

| | | |
|--|-----------|---|
| (51) International Patent Classification ⁵ : C07D 305/14, 413/12, A61K 31/335 | A1 | (11) International Publication Number: WO 94/25449 (43) International Publication Date: 10 November 1994 (10.11.94) |
| (21) International Application Number: PCT/US94/04519 (22) International Filing Date: 25 April 1994 (25.04.94) (30) Priority Data: 08/053,902 26 April 1993 (26.04.93) US (71) Applicant: HAUSER CHEMICAL RESEARCH, INC. [US/US]; 5555 Airport Boulevard, Boulder, CO 80301 (US). (72) Inventors: MURRAY, Christopher, K.; 5555 Airport Boulevard, Boulder, CO 80301 (US). BECKVERMIT, Jeffrey, T.; 5555 Airport Boulevard, Boulder, CO 80301 (US). ZIEBARTH, Timothy, D.; 5555 Airport Boulevard, Boulder, CO 80301 (US). (74) Agent: EDMUNDSON, Dean, P.; 1136 East Stuart Street, Suite 2160, Fort Collins, CO 80525 (US). | | (81) Designated States: AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i> |
| (54) Title: OXIDATION PRODUCTS OF CEPHALOMANNINE (57) Abstract Antineoplastic taxol derivatives are derived by selective oxidation of the alkene portion of the side chain of cephalomannine. The derivative displays high activity in promoting assembly of microtubulin and also displays cytotoxic activity against malignant cells. | | |

#44

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

| | | | | | |
|----|--------------------------|----|--|----|--------------------------|
| AT | Austria | GB | United Kingdom | MR | Mauritania |
| AU | Australia | GE | Georgia | MW | Malawi |
| BB | Barbados | GN | Guinea | NE | Niger |
| BE | Belgium | GR | Greece | NL | Netherlands |
| BF | Burkina Faso | HU | Hungary | NO | Norway |
| BG | Bulgaria | IE | Ireland | NZ | New Zealand |
| BJ | Benin | IT | Italy | PL | Poland |
| BR | Brazil | JP | Japan | PT | Portugal |
| BY | Belarus | KE | Kenya | RO | Romania |
| CA | Canada | KG | Kyrgyzstan | RU | Russian Federation |
| CF | Central African Republic | KP | Democratic People's Republic of Korea | SD | Sudan |
| CG | Congo | KR | Republic of Korea | SE | Sweden |
| CH | Switzerland | KZ | Kazakhstan | SI | Slovenia |
| CI | Côte d'Ivoire | LI | Liechtenstein | SK | Slovakia |
| CM | Cameroon | LK | Sri Lanka | SN | Senegal |
| CN | China | LU | Luxembourg | TD | Chad |
| CS | Czechoslovakia | LV | Latvia | TG | Togo |
| CZ | Czech Republic | MC | Monaco | TJ | Tajikistan |
| DE | Germany | MD | Republic of Moldova | TT | Trinidad and Tobago |
| DK | Denmark | MG | Madagascar | UA | Ukraine |
| ES | Spain | ML | Mali | US | United States of America |
| FI | Finland | MN | Mongolia | UZ | Uzbekistan |
| FR | France | | | VN | Viet Nam |
| GA | Gabon | | | | |

DescriptionOxidation Products of CephalomannineTechnical Field

5 This invention relates to taxane derivatives. More particularly, this invention relates to oxidation products of cephalomannine. In another aspect, this invention relates to techniques for producing taxol derivatives from cephalomannine.

Background Art

10 Taxol, 1, a material occurring in nature, and extracted from Taxus brevifolia (i.e., the Pacific yew tree) and other biomass has been identified as having significant tubulin binding (Schiff, P. B. et al., "Promotion of Microtubule Assembly in vitro by Taxol," 15 Nature, Vol. 277: 665-67 (Feb. 1979)) and, when delivered to the cell, cytotoxic activity which has been demonstrated through Phase III clinical trials. Taxol was recently approved for the treatment of refractory ovarian cancer by the Food and Drug Administration.

20 Taxotere, 2, a semisynthetic derivative of taxol with improved water solubility, has been compared with taxol in Phase I clinical trials. Taxotere is slightly more active as a promoter of tubulin polymerization, 1.5-fold more potent as an inhibitor of replication in mouse macrophage-like J774.2 cells and in P388 murine leukemia cells, and at 25 least fivefold more potent in taxol resistant tumor cells (Pazdur, R. et al., "Phase I Trial of Taxotere: Five-Day Schedule", Journal of the National Cancer Institute, 1781, (1992)). The structural differences between taxol 1 and 30 taxotere 2 are minor (Figure 1), yet enhanced in vitro tubulin binding activity is observed for taxotere.

Consequently, it is difficult to predict the relative potency of a taxol analogue for microtubulin polymerization activity based on small changes in the overall structure. An examination of Kingston's Review, (Kingston, D. G. I.,
5 "The Chemistry of Taxol", Pharmacology and Therapeutics, 52: 1-34, (1991)), provides an overall view of the complexity of the structure-activity relationship of taxol analogues. It is clear that minor structural changes can cause major changes in tubulin binding activity and
10 cytotoxicity. These changes can even completely eliminate activity. In addition, other factors such as greater water solubility and lower toxicity exist, which must be strongly considered when evaluating the efficacious nature of therapeutic agents.

15 The novel synthetic taxol derivatives described herein have not heretofore been described nor has the literature suggested that such new derivatives would exhibit tubulin assembly or advantageous cytotoxic activity.

Disclosure of Invention

20 There has now been discovered a new compound that displays in vitro tubulin binding and cytotoxic activity similar to taxol. The new antineoplastic taxol derivative is derived by selective oxidation of the alkene portion of the side chain of cephalomannine 3. The formation of this
25 new taxol derivative from cephalomannine has not been described previously and provides in high yield the new derivative.

It is an object of this invention to provide a new semisynthetic taxol derivative that displays unexpectedly
30 high activity in promoting the assembly of microtubulin in vitro and cytotoxic activity against B16 melanoma cells, for example.

It is another object of this invention to provide a pharmaceutical composition which is effective in inhibiting the growth of tumor cells.

It is a further object of this invention to provide
5 methods for producing the new taxol derivative.

Other objects and advantages of the present invention will be apparent from the following detailed description and the accompanying drawings.

Best Mode for Carrying Out the Invention

10 This invention relates to the treatment of cephalomannine 3, a close natural analogue of taxol, with a strong oxidizing agent, e.g. ozone, to generate in good yield a new derivative mixture. The cephalomannine starting material can be isolated in a conventional manner
15 such as described in a recent publication (Rao, Koppaka V., "Method for the Isolation and Purification of Taxane Derivatives", International Publication Number, WO 92/07842, May 14, 1992).

Treatment of an ether and/or hydrocarbon and/or
20 alcoholic and/or chlorinated solvent solution of cephalomannine between -78° C and room temperature, with 5 to 1000 equivalents of ozone, followed by purging with an inert gas, results in the formation of the α -keto-
amide(pyruvamide)/ α -ketal-amide derivative mixture 4a, 4b
25 (see Figure 2). The transformation is very selective for the side chain alkene and over-oxidation can be avoided, i.e., oxidation of the tetrasubstituted alkene in ring A and other functional groups can be prevented, if an amount of ozone is added which is sufficient to completely oxidize
30 the tiglate amide functional group (see R for compound 3 of Figure 1), while also avoiding oxidation elsewhere in the molecule. The correct stoichiometry is determined by calibrating the ozone generator and by monitoring the reaction using high pressure liquid chromatography (HPLC).

A description of the method for monitoring reactions by HPLC is contained in the Examples section below.

The new synthetically modified taxane derivative mixture, hereafter designated **4ab**, was characterized by spectroscopic analysis. The ¹H-NMR and ¹³C-NMR spectra of the equilibrium mixture shows both **4a** and **4b** present in a ratio of 4:1 in chloroform-d (CDCl₃). When the compounds are analyzed by ¹³C-NMR in methyl-d₃ alcohol-d (CD₃OD), the ratio changes to approximately 1:1. In CDCl₃ solvent, the ketone (6'-**4a**) carbonyl resonance at 195.4 ppm is approximately four times larger than the hemiketal (6'-**4b**) carbon resonances at 102.5 and 105.4 ppm (two diastereomers for **4b**). In CD₃OD solvent, the ketone (6'-**4a**) carbonyl resonance at 197.2 ppm is approximately the same peak height (non-quantitative ¹³C-NMR experiment) as the ketal/hemiketal carbon resonances at 97.9, 101.9, 104.2, 105.1, and 106.5 ppm. The five different ketal/hemiketal carbon resonances in CD₃OD are attributed to the diastereomeric ketal carbons represented in **4b**, and solvent addition to the open and closed forms of **4ab**. It should be emphasized here that the two forms **4a** and **4b** are rapidly interconverting in solution at room temperature. Isolation of exclusively one form from the other without resorting to chemical conversion of the mixture has not been observed.

The new synthetic taxane derivative mixture **4ab** was characterized by chemical conversion to other new taxane derivatives. The crude ozonolysis reaction mixture was treated with acetic anhydride in pyridine (see Figure 3) to generate the two compounds shown (5 and 6). The 2'-OH is acetylated in both compounds as shown so the equilibrium between open and closed forms for the starting material (**4ab**) is not possible. A similar result was observed upon silylation using triethylsilyl chloride in pyridine (see Figure 4). The triethylsilyl derivative **7** cannot cyclize because the 2'-OH is blocked. All of the compounds

described here were characterized by spectroscopic techniques (for synthesis method and characterization data see the section below titled Examples).

The new synthetic taxane derivative mixture **4ab** was synthesized by an additional method of organic chemical synthesis from a different starting material. As shown in Figure 5, the diol **8** is available via dihydroxylation of cephalomannine using established methodology (Kingston, D. G. I., et al, "Modified Taxols, 7. A Method for the Separation of Taxol and Cephalomannine", J. Nat. Prod. 55: pp 259-261, (1992)). When the diol **8** is treated with sodium periodate, the expected compound **4a** (in equilibrium with the cyclized form **4b**), is formed in very good yield. The oxidative cleavage of a vicinal diol functional group similar to the side-chain portion of **8**, is known to yield carbonyl compounds similar to the ketoamide group of **4a** (Sklarz, B., "Organic Chemistry of Periodates", Quarterly Reviews, pp 3-28, (1967)). The methodology described here is another structure proof of **4ab**. The synthesis method shown in Figure 5 provides a product mixture with identical spectral and chromatographic analyses as the product mixture from reacting ozone with cephalomannine (see Figure 2).

The cleavage of the diol functional group in compound **8** is achieved by using an effective amount of an oxidizing agent. Effective oxidizing agents include, but are not limited to, periodic acid and salts thereof, lead tetraacetate, sodium bismuthate, tetrabutylammonium periodate, manganese dioxide, pyridinium chlorochromate, and potassium permanganate. The oxidizing agents listed are not ranked according to effectiveness in performing the oxidation step. The relative effectiveness of the various possible oxidizing agents depends upon the concentration employed and other conditions of the reaction.

The new synthetic taxane derivative mixture **4ab** was also synthesized by a variation on the two-step method shown in Figure 5. Treatment of compound 3 in a two-phase solvent system as shown in Figure 6, with sodium periodate and ruthenium trichloride catalyst, results in a mixture with identical chromatographic and spectral (ultraviolet) characteristics to **4ab**. The sodium periodate/ruthenium trichloride oxidative cleavage of an internal alkene functional group similar to the side-chain portion of 3, is known to yield carbonyl compounds similar to the ketoamide group of **4a** (Carlsen, P. H. J., et al., "A Greatly Improved Procedure for Ruthenium Tetraoxide Catalyzed Oxidations of Organic Compounds", J. Org. Chem., pp 3936 - 3938, (1981)). In a similar manner, other transition metal catalysts that are capable of diol oxidation when used in combination with oxidants such as periodate or hydroperoxide, can be used for the oxidative cleavage of the alkene portion of the side chain of cephalomannine. The ruthenium trichloride/periodate oxidation methodology described here constitutes another structure proof of **4ab**, and provides a third oxidation method for the synthesis of **4ab** from cephalomannine 3.

The new compound mixture, **4ab**, shows good tubulin binding and cytotoxicity activity with in vitro testing. The in vitro test results are comparable to results for taxol. Tubulin binding and cytotoxicity data for cephalomannine and the synthetic derivatives described herein are included for comparison. The tubulin testing was done exactly as described by Himes (Georg, G. I., et al., "Synthesis of Biologically Active Taxol Analogues with Modified Phenylisoserine Side Chains", J. Med. Chem. Vol. 35: 4230, (1992)). See Table 1 for the data. Taxol has been included in Table 1 for reference. In addition, each sample is compared to taxol in the columns: ED₅₀/ED₅₀ Taxol (for Tubulin Assembly), and ED₅₀/ED₅₀ Taxol (for B16

Proliferation); taxol shows a value of approximately 1 in these columns. A number less than 1 in these columns indicates greater activity than taxol. A number greater than 1 in these columns indicates lower activity than taxol. The error in the tests appears to be $\pm 10 - 20\%$. The data clearly shows that the α -ketoamide-taxane mixture **4ab** has activity comparable to or superior to taxol in the in vitro tublin assembly and B16 Proliferation tests. These tests have been used and relied upon by experimentalists in this field to determine the potential efficacy of a taxol derivative for the treatment of cancer. The data in Table 1 also demonstrates the dramatic difference in activity between structurally similar taxane compounds; see for example, data for compounds **4ab**, **3**, and **8**.

The synthesis, characterization and in vitro test methods for the new taxol derivatives are illustrated by the following examples:

Examples

Chemistry. All solvents and reagents employed were used as received from the manufacturer except pyridine and acetic anhydride which were distilled prior to use. Reactions were monitored by thin-layer chromatography ("TLC") using 0.20 mm. E. M. Industries Silica Gel 60 (aluminum support) silica gel plates. Reactions were also monitored by high pressure liquid chromatography ("HPLC"). Aliquots of crude reaction mixtures for HPLC analysis were removed from the reaction vessel with a 3 μ l micro-pipette and diluted to 200 μ l in an HPLC sample vial (with insert). The HPLC system consists of a model L-6200 pump, Model AS-4000 or L-3000 UV/VIS/DAD detector (Hitachi Instruments, Inc.). The system was equipped with an NEC 286 computer with 40M hard drive and Lab Manager HPLC software (Hitachi Instruments, Inc.). HPLC columns used included a 4.6 mm.

X 250 mm. Phenyl column, packed with 5 μ m diphenyl material (Supelco, Inc.); a 4.6 mm. X 250 mm., 5 μ m, 60 angstrom Pentafluorophenyl (PFP) column (ES Industries); and a 4.6 mm. X 250 mm. phenyl guard column (Jones Chromatography).

5 The ozone generator used was a Polymetrics Laboratory Ozonator T-816 with an operation at 75 volts, 60 Hertz current, 5.5 psig pressure, and a flow of 2 SLMP delivering a concentration of ozone at 2.2 mg/s. The ozone flow was calibrated using the method described by the manufacturer.

10 Silica gel for flash chromatography (230 to 400 mesh) was supplied by Scientific Products. A Bruker WP-270 and ACE-300, Varian Gemini 400, and a JEOL FX90Q Spectrometer were employed for ^1H and ^{13}C NMR spectra with chemical shifts reported in ppm. relative to tetramethylsilane using

15 residual non-deuterated NMR solvent for reference. Yields refer to chromatographically pure compounds and are not optimized. Purity of products were judged to be >90 % on the basis of spectrophotometric homogeneity unless otherwise stated. Mass spectra were measured at M-Scan

20 Inc. using a VG Analytical 2-SE high field mass spectrometer. Spectroscopic analyses were determined using an Analect Diamond-20 FTIR with an XAD-Plus microscope. The instrument was equipped with an ACR Advanced Logic Research 486 computer with 200M hard drive and an Analect

25 FX80 software package.

Example 1

α -Ketoamide 4a. Cephalomannine 3 (178 mg) dissolved in CHCl_3 (5 ml) was treated with ozone (2.2 mg/s) for 90 seconds at room temperature followed by evaporation to give

30 a quantitative yield of the isomeric mixture 4ab. Resonances for the major isomer are listed. ^1H NMR (90 MHz, CDCl_3) 1.11 (s, 3H), 1.21 (s, 3H), 1.29 - 1.58 (m, 2H), 1.64 (s, 3H), 1.78 (s, 3H), 1.89 - 2.16 (m, 3H), 2.20 (s, 3H), 2.31 (s, 3H), 2.24 (s, 3H), 2.40 - 2.72 (m, 1H),

3.75 (d, $J = 6.8$ Hz, 2H), 3.98 - 4.47 (m, 3H), 4.64 (m, 1H), 4.89 (d, $J = 8.5$ Hz, 1H), 5.13 - 5.68 (m, 2H), 5.98 - 6.20 (m, 1H), 6.25 (s, 1H), 7.27 - 7.73 (m, 8H), 7.85 (d, $J = 9.4$ Hz, 1H), 8.07 (d, $J = 7.7$ Hz, 2H). ^{13}C NMR (12 MHz, CDCl_3) 9.54, 14.65, 20.76, 21.65, 22.51, 24.37, 26.79, 35.62, 35.62, 43.12, 45.72, 55.00, 58.48, 72.04, 72.04, 73.31, 74.96, 75.58, 76.44, 79.00, 81.16, 84.32, 126.97, 126.97, 128.60, 128.60, 128.60, 128.89, 128.89, 129.16, 130.11, 130.11, 133.24, 133.65, 137.14, 141.68, 159.70, 166.88, 170.30, 171.07, 171.93, 195.94, 203.55. The diagnostic signals in the ^{13}C -NMR spectrum for the minor isomer in CDCl_3 (carbon 6' of the two diastereomers of **4b**) are 102.52 and 105.35 ppm. The diagnostic signals in CD_3OD for **4ab**, including solvent addition (CD_3OD) to both **4a** and **4b** are 97.9, 101.9, 104.2, 105.1, and 197.2 ppm for carbon 6'. FTIR (neat, cm^{-1}) 981.6 (m), 1025.9 (m), 1070.3 (m), 1108.9 (m), 1178.3 (m), 1241.9 (s), 1373.1 (m), 1724.0 (s), 2900.4 (w), 2940.9 (w), 3064.3 (w), 3413.4 (m), 3490.5 (m). Mass Spectrum (FAB, glycerol/thioglycerol matrix) m/z 821 ($M + 1$) $^+$.

Example 2

2'-Acetyl- α -ketoamide 5 and 2',7-bis(acetyl)- α -ketoamide 6.

Cephalomannine **3** (320 mg) in CH_2Cl_2 (4 ml) was treated with ozone (2.2 mg/s) for 205 seconds, purged with nitrogen, and evaporated to dryness. The oxidized cephalomannine in CH_2Cl_2 (1.5 ml) was cooled to 0°C , acetic anhydride (0.145 ml) and pyridine (0.156 ml) were added sequentially. The reaction was stirred at 0°C for 2 hours followed by stirring for an additional 21 hours at room temperature. After diluting with methylene chloride the mixture was washed with 3N HCl (3x), saturated NaHCO_3 , and brine. The solution was dried over MgSO_4 , and evaporated. Flash chromatography on silica gel (45/55, 55/45, 75/25

ethyl acetate/hexane) afforded 2 products. The first was 191 mg (58%, white, R_f = 0.16 50/50 ethyl acetate/hexane) corresponding to 5. ^1H NMR (270 MHz, CDCl_3) 1.10 (s, 3H), 1.21 (s, 3H), 1.61 (s, 3H), 1.81 (s, 1H), 1.86 (d, J = 1.2 Hz, 3H), 2.10 (s, 1H), 2.12 (s, 3H), 2.19 (s, 3H), 2.35 (s, 3H), 2.38 (s, 3H), 1.68 - 2.58 (m, 4H), 3.76 (d, J = 7.0 Hz, 1H), 4.13 (d, J = 8.2 Hz, 1H), 4.27 (d, J = 8.2 Hz, 1H), 4.35 - 4.46 (m, 1H), 4.95 (dd, J = 1.9, 7.2 Hz, 1H), 5.35 (d, J = 4.1 Hz, 1H), 5.57 (dd, J = 3.5, 9.7 Hz, 1H), 5.63 (d, J = 7.0 Hz, 1H), 6.14 (t, J = 9.4 Hz, 1H), 6.24 (s, 1H), 7.25 - 7.65 (m, 8H), 7.70 (d, J = 9.4 Hz, 1H), 8.11 (dd, J = 1.8, 7.0 Hz, 2H). ^{13}C NMR (68 MHz, CDCl_3) 9.78, 14.86, 20.56, 20.99, 22.30, 22.93, 24.60, 26.95, 35.90, 35.90, 43.52, 46.06, 53.39, 58.76, 72.29, 72.49, 74.28, 75.47, 75.90, 76.62, 79.59, 81.31, 84.65, 127.11, 127.11, 129.04, 129.04, 129.22, 129.38, 129.38, 129.70, 130.49, 130.49, 133.19, 134.05, 136.64, 143.04, 159.98, 167.20, 168.21, 170.13, 170.22, 171.62, 196.49, 204.09. FTIR (neat, cm^{-1}) 710 (w), 934 (m), 1026 (m), 1070 (s), 1242 (s), 1271 (s), 1373 (s), 1728 (s), 2941 (m), 2960 (m), 3410 (w), 3514 (m). Mass spectra (FAB, m-nitro benzyl alcohol matrix) m/z 863 ($M + 1$) $^+$. The second was 98 mg (28%, white, R_f = 0.34 50/50 ethyl acetate/hexane) corresponding to 6. ^1H NMR (270 MHz, CDCl_3) 1.13 (s, 3H), 1.19 (s, 3H), 1.75 (s, 3H), 1.79 (s, 1H), 1.88 (d, J = 1.2 Hz, 3H), 1.96 (s, 3H), 2.13 (s, 6H), 1.61 - 2.34 (m, 3H), 2.37 (s, 3H), 2.38 (s, 3H), 2.47 - 2.61 (m, 1H), 3.88 (d, J = 7.0 Hz, 1H), 4.12 (d, J = 8.8 Hz, 1H), 4.29 (d, J = 8.2 Hz, 1H), 4.94 (d, J = 8.2 Hz, 1H), 5.38 (d, J = 3.5 Hz, 1H), 5.46 - 5.59 (m, 2H), 5.63 (d, J = 7.0 Hz, 1H), 6.11 (t, J = 9.1 Hz, 1H), 6.18 (s, 1H), 7.28 - 7.65 (m, 8H), 7.71 (d, J = 9.4 Hz, 1H), 8.11 (dd, J = 1.8, 7.0 Hz, 2H). ^{13}C NMR (68 MHz, CD_2Cl_2) 11.05, 14.48, 20.57, 20.86, 21.17, 21.46, 22.88, 24.58, 26.58, 33.58, 35.72, 43.59, 47.44, 53.46, 56.24, 71.77, 72.24, 74.16, 74.91, 75.45, 76.53, 79.17,

81.22, 84.19, 127.13, 127.13, 129.04, 129.04, 129.22, 129.36, 129.36, 129.65, 130.47, 130.47, 133.03, 134.09, 136.69, 141.27, 167.13, 168.32, 168.32, 169.27, 170.06, 170.10, 170.10, 170.46, 202.19. FTIR (neat, cm^{-1}) 710 (w), 1049 (m), 1068 (m), 1240 (s), 1269 (s), 1697 (m), 1728 (s), 1751 (s), 2956 (w), 3410 (w), 2523 (bw). Mass spectrum (FAB, m-nitro benzyl alcohol matrix) m/z 905 ($M + 1$)⁺.

Example 3

2',7-Bis(triethylsilyl)- α -ketoamide 7.

To α -ketoamide 4a (75.5 mg) dissolved in pyridine (4.6 ml) was added triethylsilyl chloride (0.31 ml). The reaction mixture was mixed at room temperature for 22 hours followed by CH_2Cl_2 dilution. The organic phase was washed sequentially with 3N HCl (2x), saturated NaHCO_3 , and brine. It was then dried over MgSO_4 and evaporated to a solid. Flash chromatography on silica gel (25/75 ethyl acetate/hexane) afforded 39 mg (41%) of a white solid (R_f = 0.24, 25/75 ethyl acetate/hexane). ^1H NMR (300 MHz, CD_2Cl_2) 0.35 - 0.63 (m, 12H), 0.77 - 0.97 (m, 18H), 1.21 (s, 6H), 1.66 (s, 3H), 1.94 (d, J = 1.1 Hz, 3H), 1.81 - 1.91 (m, 1H), 1.99 - 2.20 (m, 1H), 2.15 (s, 3H), 2.35 (s, 3H), 2.32 - 2.42 (m, 1H), 2.50 (s, 3H), 2.51 - 2.58 (m, 1H), 3.81 (d, J = 7.1 Hz, 1H), 4.15 (d, J = 8.2 Hz, 1H), 4.29 (d, J = 8.4 Hz, 1H), 4.47 (dd, J = 6.7, 10.6 Hz, 1H), 4.63 (d, J = 2.7 Hz, 1H), 4.95 (m, 1H), 5.40 (dd, J = 2.7, 9.4 Hz, 1H), 5.67 (d, J = 7.2 Hz, 1H), 6.18 (t, J = 9.2 Hz, 1H), 6.42 (s, 1H), 7.27 - 7.44 (m, 5H), 7.49 - 7.67 (m, 3H), 7.79 (d, J = 9.4 Hz, 1H), 8.15 (dd, J = 1.5, 7.0 Hz, 2H). ^{13}C NMR (75 MHz, CD_2Cl_2) 4.68, 4.68, 4.68, 5.59, 5.59, 5.59, 6.66, 6.66, 6.66, 6.89, 6.89, 6.89, 10.34, 14.44, 21.01, 21.73, 23.18, 24.58, 26.69, 35.85, 37.57, 43.67, 47.12, 56.01, 58.65, 71.99, 72.67, 75.22, 75.34, 75.43, 76.74, 79.38, 81.36, 84.45, 127.06, 127.06, 128.61, 128.61, 129.04, 129.09, 129.09, 129.09, 129.79, 130.51, 130.51, 134.00, 138.20,

140.53, 160.01, 167.27, 169.44, 170.36, 171.65, 196.76, 201.95. FTIR (neat, cm^{-1}) 733 (w), 746 (w), 1003 (m), 1018 (m), 1109 (s), 1138 (m), 1242 (s), 1267 (s), 1369 (m), 1695 (m), 1726 (s), 2877 (m), 2914 (m), 2958 (m), 3028 (w), 3404 (w). Mass Spectrum (FAB, m-nitro benzyl alcohol matrix) m/z 1049 ($M + 1$)⁺.

Example 4

Cephalomannine diol 8. Cephalomannine, 3, was oxidized (stoichiometric reaction) as described by Kingston (Kingston, D. G. I., et al, Modified Taxols, 7. A Method For The Separation Of Taxol And Cephalomannine", Journal of Natural Products, Vol. 55, 259-261, (1992)). The spectrophotometric analyses correspond to Kingston's reported values.

Example 5

α -Ketoamide 4a from diol 8. The diol 8 (79 mg) was dissolved in THF (0.400 ml), and water (0.342 ml) and NaIO_4 (59 mg) were added. After stirring for five minutes a white precipitate appeared, and after 20 hours reaction time, analysis by HPLC showed the reaction was complete. It was evaporated on a rotary evaporator, reconstituted with EtOAc/water and separated. The aqueous layer was extracted again with EtOAc and the combined organics were washed with sat. Na_2SO_3 and brine. The mixture was dried over MgSO_4 and evaporated to yield 62 mg (83%) of a white solid. The data (^1H -NMR, ^{13}C -NMR, IR and FAB-MS) for this sample matched exactly the data for compound 4ab prepared by ozonolysis of cephalomannine (see Example 1).

Example 6

α -Ketoamide 4a from cephalomannine 3 via RuCl_4 /periodate oxidation. Cephalomannine 3 (24.6 mg) was dissolved in carbon tetrachloride (0.06 ml), acetonitrile

(0.06 ml), and water (0.092 ml). To this biphasic mixture was added NaIO_4 (26.3 mg; 4.1 equivalents) and ruthenium trichloride (0.15 mg; 2.2 mol%) were added. After stirring for five minutes a red/brown precipitate appeared, and
5 after 1 hour the reaction was stopped. It was worked up with methylene chloride, washed with brine, and dried over anhydrous MgSO_4 . After concentrating to a light yellow solid the sample was analyzed by HPLC. The retention time, peak shape and UV spectrum of the sample was compared with
10 a previously prepared sample of **4ab**. The data for this sample matched exactly the data for compound **4ab** prepared by ozonolysis of cephalomannine (see Example 1).

Example 7

Biological Testing. B16 Melanoma Cell Proliferation.

15 Cells were seeded in 24-well plates at 7.5×10^4 cells/well and grown in Dulbecco's modified minimal essential medium (MEM) containing 10% bovine calf serum at 37° C for 24 hours in a 97% humidified atmosphere of 5.5% CO_2 . The medium was then replaced with fresh medium containing taxol
20 or its derivatives and dissolved in DMSO in concentrations ranging from 7.5×10^{-9} M to 1×10^{-7} M for taxol and other derivatives. The final concentration of DMSO in the cell medium was 0.5% or less. This amount of DMSO did not have any effect on cell proliferation as determined from control
25 experiments. After 40 hours, the cells were released by trypsinization and counted in a Coulter counter.

Tubulin Preparation and Assembly. Tubulin free of microtubule-associated proteins was purified from bovine brain as previously described (Algaier, J.; Himes, R. H.,
30 "The Effect of Dimethyl Sulfoxide on the Kinetics of Tubulin Assembly" Biochim. Biophys. Acta, Vol 954, pp 235 - 243, 1988). The assembly reaction was done at 37° C in PEM buffer (0.1 M Pipes, pH 6.9, 1 mM EGTA, and 1 mM MgSO_4) at a protein concentration of 1 mg/ml (10 μM) in the presence

of taxol or taxol analogues and 0.5 mM GTP. The reaction was monitored by the increase in the apparent absorbance at 350 nm.

TABLE 1

| Compound ^a | Tubulin Assembly ^b | | B16 Proliferation ^c | |
|-----------------------|-------------------------------|---|--------------------------------|---|
| | ED ₅₀ ^d | ED ₅₀ /ED ₅₀ Taxol | ED ₅₀ ^e | ED ₅₀ /ED ₅₀ Taxol |
| 1 | 1.08 | 1.26 | 21.4 | 0.95 |
| 4ab | 1.24 | 1.46 ^g | 17.1 | 0.75 ^g |
| 3 | 0.70 | 0.82 ^h | 33.5 | 1.49 ^h |
| 8 | 3.3 | 3.4 ⁱ | >854 | >38 ^{ij} |
| 5 | >8.54 ^f | >10 ^h | >854 ^f | >38 ^h |
| 6 | >8.54 ^f | >10 ^h | >854 ^f | >38 ^h |
| 7 | >8.54 ^f | >10 ^h | >8540 ^f | >380 ^h |

^aMethanol (0.5 ml) was added to each vial. Concentrations were determined from the extinction coefficients (absorbance is of a 1% wt./vol. (mg/ml) solution in methanol at 227 nm).

^bTubulin at 1 mg/ml was incubated with various concentrations of the compounds at 37°C for 15 minutes in 0.5 ml of PEM buffer (0.1 M Pipes, 1 mM EGTA, 1 mM MgSO₄, pH 6.9). Samples were centrifuged and the protein concentration on the supernatant was determined.

^cB16 Melanoma cells were incubated with various concentrations of the compounds for about 40h at 37°C.

^dThe concentration in ng/ml which reduces the supernatant protein concentration by 50%.

^eThe concentration in ng/ml which reduces the number of cells by 50% compared to a control.

^fThe highest concentration used without achieving 50% inhibition.

^gED₅₀ for taxol in the assembly assay was 0.85 µg/ml. In the B16 assay it was 22.7 ng/ml.

^hED₅₀ for taxol in the assembly assay was 0.854 µg/ml. In the B16 assay it was 22.5 ng/ml.


ⁱED₅₀ for taxol in the assembly assay was 0.97 µg/ml. In the B16 assay it was 22.7 ng/ml.

- 5 ^jThe highest concentration used was 854 ng/ml without achieving 50% inhibition.

16

Claims


1. An antineoplastic derivative of taxol having the following formula:



4a

2. A pharmaceutical composition comprising an effective amount of the antineoplastic derivative of claim 1 as an active ingredient, and a pharmaceutically-acceptable carrier.


3. A cytotoxic composition comprising:



4a

17

4. A method for inhibiting growth of cancer cells comprising contacting said cells with an effective amount of a composition comprising:



4a

5. A method for oxidizing cephalomannine comprising contacting cephalomannine with ozone.

6. A method for oxidizing cephalomannine to produce the compound of claim 1, the method comprising the steps of:


- (a) oxidizing cephalomannine with osmium tetroxide to produce a diol derivative; and
- (b) contacting said diol derivative with a diol cleaving agent to yield the compound of claim 1.

7. A method in accordance with claim 6, wherein said diol cleaving agent comprises periodate.

8. A method for oxidizing cephalomannine to produce the compound of claim 1, the method comprising contacting cephalomannine with a transition metal catalyst capable of alkene oxidation in the presence of periodate or hydroperoxide.


18

9. An antineoplastic derivative of taxol having the following formula:




4a

in equilibrium with



4b

FIG. 1



- 1 $R = C_6H_5CO$, $R' = Ac$ taxol
- 2 $R = C_4H_9OCO$, $R' = H$ taxotere
- 3 $R = CH_3CH=CCO$, $R' = Ac$ cephalomannine
 $\quad \quad \quad |$
 $\quad \quad \quad CH_3$

2/6




FIG. 2






FIG. 3

FIG. 4



5/6



6/6




FIG. 6

INTERNATIONAL SEARCH REPORT

International Application No.
PC1/US 94/04519

A. CLASSIFICATION OF SUBJECT MATTER
IPC 5 C07D305/14 C07D413/12 A61K31/335

According to International Patent Classification (IPC) or to both national classification and IPC.

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 5 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

| Category * | Citation of document, with indication, where appropriate, of the relevant passages | Relevant to claim No. |
|------------|--|-----------------------|
| A | JOURNAL OF MEDICINAL CHEMISTRY vol. 35 , 1992 , WASHINGTON US pages 4230 - 4237 GUNDA GEORG 'SYNTHESIS OF BIOLOGICALLY ACTIVE TAXOL ANALOGUES WITH MODIFIED PHENYLISOSERINE SIDE CHAINS.' cited in the application see page 4230 - page 4231 ----- | 1,2 |

☐ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

23 August 1994

Date of mailing of the international search report

- 5. 09. 94

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+ 31-70) 340-3016

Authorized officer

Francois, J